

Refining the Drag Parameter for Coronal Mass Ejection propagation models

D. Del Moro⁽¹⁾, F. Berrilli⁽¹⁾, R. Foldes⁽¹⁾, L. Giovannelli⁽¹⁾, G. Napoletano⁽²⁾, E. Pietropaolo⁽²⁾

⁽¹⁾Università degli Studi di Roma "Tor Vergata", Via della Ricerca Scientifica, 1, 00133 Rome, Italy;

⁽²⁾Università de L'Aquila, Via Vetoio, 48, 67100 Coppito L'Aquila, Italy.


INTRO:

ICME (Interplanetary Coronal Mass Ejection) are violent phenomena of solar activity that affect the whole heliosphere and the prediction of their impact on different solar system bodies is one of the primary goals of the planetary space weather forecasting.


What: an exploratory work on how the drag parameter depends on the physical parameters of the ICME, to verify and extend the work by Vršnak et al. (2010).

Scope: a more robust definition of the PDF of the drag parameter that could be used in the various approaches to ICME propagation forecasting.

Methods for CME detection and characterization




Predictions of the arrival of ICMEs in geospace are produced through use of **CME geometric models** combined with ICME propagation models, **constraining** these models with the **available Coronagraph and Heliospheric Imager** data.



Often, geometric models provide inconsistent results because assumptions inherent to each model are being invalidated

Models for ICME propagation



Predictions of the arrival of ICMEs in geospace are produced through use of CME geometric models combined with **ICME propagation models**, constraining these models with the available Coronagraph and Heliospheric Imager data.

The DBM equation and assumptions

The **Drag Based Model** approach is based on a simple equation for the ICME acceleration as

$$\frac{d^2r}{dt^2} = -\gamma \left(\frac{dr}{dt} - w \right) \left| \frac{dr}{dt} - w \right|$$


where w is the ambient solar-wind speed and γ is the so-called drag parameter (Vršnak et al., 2013).

In this framework, γ depends on the ICME mass and cross-section, on the solar-wind density and, to a lesser degree, on other parameters.

The typical working hypothesis for DBM implies that both γ and w are constant far from the Sun.

To run the codes, forecasters use empirical input values for γ and w .




- Self-Similar expansion
- Evolution entirely governed by fluid dynamics (i.e. the interplanetary magnetic field plays no role)
- Constant solar wind speed and drag parameter



The P-DBM : the PDFs.

In the DBM 'Ensemble' approaches (Dumbovich et al., 2018; Napoletano et al. 2018), the uncertainty about the actual values of such inputs are rendered by **Probability Distribution Functions** (PDFs), accounting for the values variability and our lack of knowledge.


Among those PDFs, that of γ is poorly defined due to the relatively scarce statistics of recorded values.



Our works: stats from R&C and our algorithms


Taking advantage of the large data-set of ICMEs registered by Richardson & Cane (2010), we computed how γ depends on SW velocity...

$$\pm\gamma = \frac{(v_0 - v_1)}{(v_1 - w)(v_0 - w)T}$$




The hypothesis: $\gamma(\Delta v) \rightarrow$ Simulations

...found evidence of a variation in the γ PDF if the SW is accelerating or braking the ICME propagation. By using a set of simulations of an ICME structure into the SW fluid, we try to link this change to the structure of the ICME.



The hypothesis: $\gamma(\Delta v) \rightarrow$ Simulations

...found evidence of a variation in the γ PDF if the SW is accelerating or braking the ICME propagation.
By using a set of simulations of an ICME structure into the SW fluid, we try to link this change to the structure of the ICME.



ICME simulation results:

...found evidence of a variation in the γ PDF if the SW is accelerating or braking the ICME propagation.
By using a set of simulations of an ICME structure into the SW fluid, we try to link this change to the structure of the ICME.

#	Shape	C_d^F	C_d^B	$\gamma^F (\times 10^{-7} \text{km}^{-1})$	$\gamma^B (\times 10^{-7} \text{km}^{-1})$
1		1.3	-	0.25 ÷ 2.5	-
2		2.0	-	0.4 ÷ 4	-
3		1.7	-	0.35 ÷ 3.5	-
4		1.0	1.5	0.2 ÷ 2	0.3 ÷ 3
5		2.2	3.6	0.45 ÷ 4.5	0.7 ÷ 7

Conclusions:

A dependence of γ on Δv is possible/likely, but we are at the very limit of what can be obtained by available data.

We are striving to refine the dataset to get further statistics

We are working on upgrading the simulation framework.

“The remote sensing package on Solar Orbiter, in particular the **METIS** and SoloHI instruments, may well provide the CME observations needed to improve our understanding of the time dependence of these parameters.”

Barnard et al. 2017



EXTRA SLIDES:

CME expected at 1AU at: 2018-08-25 20:00:00 UT

